

HSP90 INTERACTORS

Chaperones and relatives

- Aha1 and its homolog Hch1
- Cdc37 (p50) and its relative Hsc70 (= Cdc37L1)
- p23 (=Sba1)
- proteins with TPR motifs: Hop (=Sti1), FKBP52 (=FKBP4; and high MW plant homologs), FKBP51 (=FKBP5), FKBP8 (=FKBP38), FKBP36 (=FKBP6), *Plasmodium* FKBP35, plant TWD1, cyclophilin-40 (Cpr6 and Cpr7), PP5 (and yeast Ppt1), Tom70, probably also related Tom71=Tom72, XAP-2 (=AIP=ARA9), Cns1 and its *Drosophila* and human relatives Dpit47 and TTC4, CHIP, UNC45A (GC-UNC45) and UNC45B, She4, DnaJC7 (=Tpr2=mDj11=CCRP), CRN, WISp39 (=FKBPL), Tah1 (=Spaghetti=RPAP3), Spag1, NASP, Toc64 and OM64, TPR1 (=Ttc1), SGT (=αSGT=SGTA), DYX1C1, AtTPR1, AtTPR2, AtTPR7, AIPL1, Tom34, *Tetrahymena* Coi12p
- CS-containing p23 relatives AARSD1, SGT1 (=SUGT1), RAR1, Siah-1-interacting protein (SIP), Chp1/CHORDC1/Morgana, B-ind1, melusin, NudC and NudCL2 (=Nudcd2)
- Cereblon
- FNIP1, FNIP2
- GIGANTEA
- Hsc70/Hsp70/Hsp72/DnaK
- Hsp60
- Ids2
- Other J-domain proteins: *E. coli* CbpA and DnaJ, human DnaJ homolog Hsj1b, cyanobacterial DnaJ2, yeast Ydj1
- mtHsp70/Grp75/mortalin
- PhLP2A
- Pih1 (=Nop17) (mostly through Tah1)
- Prefoldin 4 and 6

- S100A1
- Sse1, Sse2
- Tel2-Tti1-Tti2 complex
- TIMP2
- TPR15, TPR16
- Toxoplasma Sis1-like
- Tsc1
- USP19
- USP21
- valosin-containing protein (VCP)/p97
- ZMYND10

Transcription factors

- 12(S)-HETE receptor
- AF9/MLLT3
- all vertebrate steroid receptors (GR, MR, ERα, ERβ, PR, AR)
- AGL24
- ATF3
- BBX
- BCL-6
- Bclaf1
- BES1
- BrZ7
- BZR1
- C20orf194
- CAR
- C/EBPβ
- CEBPE (C/EBPε)
- Cwt1
- CXXC1
- cytoplasmic v-erbA
- DLX6
- DMRTA1
- E1A
- E2F1 and E2F2
- EcR
- ELF3
- FOXD4L6
- FOXM1
- FOXP2
- GTF2IRD2
- Hap1
- HCFC1
- HMGA1, HMGA2
- HNF4A
- HP1BP3
- HSF1
- HSF2
- HsfA1, HsfA2, HsfB1
- IRF2
- IRF3
- ISX
- LFY
- MAFG
- Mal63
- MalR
- MAX
- Met1
- MeSRS1
- MeWRKY20
- MKX
- mod(mdg4)
- c-Myc
- Nanog
- NFIC
- NFRKB
- Notch1 (ICN1)
- NR1H3
- NR1I2
- Nrf2
- Oct4
- p53
- p73
- PAS family members: Dioxin receptor (=AhR), Sim, HIF-1α, HIF-2α, HIF-3α
- PCGF6
- PER2
- PHOX2B
- POGK
- PPARα, PPARβ, PPARγ, PPARδ
- PRDM1
- PREB
- PXR
- REST
- REV-ERBα
- RlmA (of *Aspergillus*)
- SETDB1
- SIM2
- SLFN11
- SOX1
- SOX11
- Sp1
- SREBF1
- SREBP1
- SREBP2
- STAT1
- STAT2
- STAT3 (also in caveolin-1 complexes in rafts)
- STAT5
- SUP
- TADA2A
- TBX22
- TCF25
- TDP-43

- TEAD2
- TFDP3
- THAP4
- TonEBP/OREBP
- TR β
- TRIM32
- Tup1
- Twist1
- Upc2
- Ure2
- USP1
- VDR
- VP16
- water mold *Achlya* steroid (antheridiol) receptor
- WT1
- ZBED4
- ZBTB7A
- ZBTB17
- ZBTB20
- ZC3H7B
- ZNF215
- ZNF509
- ZNF74

Kinases

- ACVR1B
- ACVR1C
- ACVR2B
- Akt/PKB
- AKT2
- ALK
- ALK1, ALK5
- ALPK1
- AMHR2
- AMPK α , AMPK γ
- ARAF
- ASK1
- ATM
- AURKC
- Aurora B
- AXL
- Bcr-Abl
- BCR-FGFR1
- BGLF4 of EBV
- BLK
- BMPR1A
- BMX
- BTK
- c-Abl
- c-Kit
- c-Mos
- CAMK1G
- CAMK2A
- CAMK2B
- CAMK2D
- CAMK2G
- CAMK4
- CAMKK1

- CAMKK2
- CAMKV
- casein kinase II α catalytic subunit
- Cdc2 (=Cdk1)
- CDK11B
- CDK14
- CDK15
- CDK18
- Cdk2, Cdk4, Cdk6, Cdk9, Cdk11
- CDK3
- CheA (E. coli)
- Chk1
- Cla4
- CLK2
- CLK3
- Cot = Tpl-2
- CSF1R
- CSNK1A1
- DCLK2
- DDR1
- DDR2
- Death-associated kinases DAPK, DAPK2, DAPK3
- DLK
- DMPK
- DYRK1B
- DYRK2
- DYRK3
- DYRK4
- eEF-2 kinase
- EGF receptor (mutant and wt)
- eIF2- α kinases HRI, Gcn2, Perk, PKR
- Eml4-Alk
- EPHA1
- EphA2
- EPHA4
- EPHB1
- EPHB6
- ErbB2
- ERBB3
- ERBB4
- ERK5
- FASTK
- FGFR1
- FGFR3-TACC3
- FGFR3 and FGFR4
- Flt3
- FLT4
- FOP2-FGFR1
- FRK
- Fused
- FYN
- Gal1
- GRK2 and GRK6
- GRK4
- GRK7
- GSK3A

- GSK3 β
- HCK
- HER3
- HIPK2
- HIPK4
- HopBF1 effectors
- ICK
- INSRR
- Insulin receptor
- Insulin-like growth factor 1 receptor
- Integrin-linked kinase
- IP6K2
- IRAK-1
- IRAK2
- IRAK3
- Ire1 α
- ITK
- I κ B kinases (IKK) α , β , γ , ϵ
- JAK1
- JNK
- KSR
- LATS1, LATS2
- LCK
- LIMK1
- LIMK2
- Lkb1
- LMTK3
- LRRK2
- LYN
- MAP2K5
- MAP2K7
- MAP3K12
- MAP3K15
- MAP3K2
- MAP3K6
- MAP3K9
- MAP4K1
- MAP4K2
- MAP4K4
- MAPKKK (MEKK) YODA
- MAPK15
- MAPK4
- MAPK6
- MAPK7
- MAST1
- MAST2
- MATK
- MEK
- MEKK1 and MEKK3
- MERTK
- MET
- Mik1
- MINK1
- MLK3
- MLKL
- MOK, MAK, MRK
- MpkA (of *Aspergillus*)
- Mps1 (=TTK)
- mTOR
- MUSK

- MYLK2
- MYLK3
- MYLK4
- NEK11
- NEK8
- NEK9
- NIK
- NME1
- NPM-AIk
- NPR2
- NTRK1
- NTRK2
- NTRK3
- NUAK2
- Nucleophosmin-Anaplastic Lymphoma Kinase
- p38
- p90RSK
- PAK6
- PASK
- Pbs2
- PDGFRB
- PDIK1L
- PDK1
- PGK1
- PI4K2A
- PI4KII β
- Pim-1
- PIM2
- PIM3
- Pink1
- PKC λ , PKC ϵ and other PKCs
- PKM2
- PKN1
- PKN2
- platelet-derived growth factor receptor α
- Pik1
- Pik3
- Pnck
- pp60v-src, c-src
- PRKAA2
- PRKACB
- PRKCA
- PRKCB
- PRKCG
- PRKCH
- PRKCI
- PRKQC
- PRKCZ
- PRKD1
- PRKD2
- PRKD3 (PKD3)
- PRKDC
- PRKG2
- PRKX
- PRKY
- PSKH1
- PSKH2
- PTK2
- PTK2B

- PTK6
- PTK6
- Raf-1, B-Raf, Ste11
- RET
- RET/PTC1
- RIP1
- RIP3
- Ron
- ROR1
- ROR2
- RPS6KA1
- RPS6KA2
- RPS6KA3
- RPS6KA5
- RPS6KA6
- RPS6KB1
- RPS6KC1
- RPS6KL1
- Ryk
- SGK-1
- SGK2
- SGK223
- SGK3
- Slit2
- src related tyrosine kinases: fer, fes, fgr, fps, lck, yes
- SRPK1
- SRPK3
- SSCMK1
- STK32B
- STK32C
- STK33
- STK38
- STK38L
- STYK1
- SYK
- TAK1
- TAOK3
- TBK1
- TESK1
- TESK2
- TGF β receptors I and II
- TIE1
- TNK1
- TNK2
- TNNI3K
- TP53RK
- TrkA I and III
- TrkB
- TSSK1B
- TSSK2
- TSSK3
- TSSK4
- TSSK6
- Tyk2
- TYRO3
- Ulk1/Atg1
- VEGFR1, VEGFR2
- Wee1, Swe1
- WNK4
- ZAP-70

Others

<ul style="list-style-type: none"> - 2Q2 - Act1 (=TRAF3IP2) - Acsl4 - Adenosine A_{2A} receptor - α_{2C} adrenergic receptor - Ago3 - AHH1 - AID - AIP56 - Aldo-keto reductase 1B10 - ANAPC2 - Anillin - ANKMY2 - Annexin A2 - ANP32C/D - Apaf-1 - APEX2 - APH1 - APLP2 - Apn2 - apoB - APOBEC-3B, -3C, -3G - APRF1 - Arb1 - ARD1 - Argonaute-1 (Ago1) - Argonaute-2 (=Ago2=GERp95) - Argonaute-4 (Ago4) - ARMC5 - ArtAB - ASB17 - ASB2 - ASB3 - ASB4 - ASB6 - ASL - ASS1 - ATG8 (GABARAP) proteins - ATG8b, ATG8c, ATG8e, ATG12 - Atp3 - Axin 1 - BAK1 - BALF5 of EBV - BCAP (PIK3AP1) - Bcl-2 - Bcl-xL - Beclin 1 - Bid - BIN2 - BLM helicase - Bms1 - BPIFB4 - BRAT1 - BRCA1 - BRCA2 - BRD4 - BRI1
--

- | | | |
|---|---|---|
| - BRMS1 | - CYP2E1 | - FBXO9 |
| - BTRC | - CYP3A4 | - FBXW11 |
| - c-IAP1 | - Cyr1 | - FBXW2 |
| - calcineurin (Cna2; catalytic subunit) | - cytoskeletal proteins: actin, tubulin (including ciliary β 4-tubulin), myosin (including Myo3B) | - FBXW5 |
| - calmodulin | - DBC2 | - FBXW7 |
| - calmodulin methyltransferase | - DDX5 | - FGAMS |
| - calpain-1 | - Ded1 | - FHIT |
| - calponin | - DEDD | - Fibronectin |
| - CARM1 | - Dengue virus protein E | - Filamin A |
| - Caspase-8 | - Dengue virus proteins NS1/2B/3/4B/5 | - FliN, FliI (E. coli) |
| - β -catenin | - DET1 | - FLIP _S and FLIP _L |
| - CB2 cannabinoid receptor | - Dicer 1 | - Folliculin |
| - Ccp1 | - Diphtheria toxin A | - free $\beta\gamma$ subunit of G protein |
| - CCDC117 | - DNA helicase Ssl2 | - FtsZ |
| - CD38 type III | - DNA polymerase α | - G2E3 |
| - CD79a | - DNA polymerase λ | - GAN |
| - CD80 (B7-1) | - DNA polymerase η | - GBP1 |
| - CD147 (Basigin) | - DnaA (E. coli) | - GC-A (ANP receptor) |
| - Cdc13 | - DNMT1 | - GC-C |
| - Cdc14 | - Drp1 | - GF14-4 |
| - Cdc25a and Cdc25c | - Dsn1 | - Gln1 |
| - Cdk5 activator p35 | - DTX4 | - GLT-1 |
| - CFTR (nascent and mutant polypeptide) | - E6 ^A E7 | - GluR1 |
| - ChAT | - EBAX-1 | - glutathione S-transferase subunit 3 (KS type) |
| - CheZ (E. coli) | - EEF1A2 | - GPX4 |
| - Chl1 | - Emc2 | - Guanylate cyclase, soluble |
| - CHMP4B | - ENC1 | - $G\alpha_0$, $G\alpha_{12}$ |
| - Chronophin | - eNOS, nNOS (?) | - Glucocerebrosidase |
| - Cineole synthase 1 | - EPRS | - GREB1 |
| - Clathrin heavy chain | - ether-a-gogo-related potassium channel (ERG = HERG = KCNH2) | - HAX-1 |
| - CLC-1 chloride channel | - EZH2 | - HDAC1 |
| - CLC-2 chloride channel | - F1F0-ATP synthase | - HDAC6 |
| - Clostridium toxin CDT | - FANCA | - HDAC11 |
| - Clostridium toxin iota | - FBXL12 | - HECTD3 |
| - Clusterin | - FBXL13 | - Hepatitis B virus core protein |
| - CND1 (Arabidopsis gene <i>At1g32730</i>) | - FBXL14 | - Hepatitis C virus protein NS3 |
| - COG complex | - FBXL15 | - Hepatitis E virus capsid protein |
| - COI1 | - FBXL18 | - HERC4 |
| - Collagen-1 | - FBXL2 | - HERC6 |
| - Complement C9 | - FBXL3 | - Histones H1, H2A, H2B, H3 and H4 |
| - COX-2 | - FBXL6 | - HMGR |
| - CPEB1, CPEB2, CPEB3 | - FBXL8 | - Hsp27 |
| - Cry1 prototoxins | - FBXL8 | - Humanin |
| - CTA1 = CtxA1 | - FBXO10 | - Huntingtin |
| - Ctf13/Skp1 component of CBF3 | - FBXO17 | - IDH1 |
| - CTNNAL1 | - FBXO18 | - IDO1 |
| - CUL1 | - FBXO24 | - Importin 4 (IPO4) |
| - CUL2 | - FBXO25 | - Importin β 1 |
| - CUL3 | - FBXO27 | - Importin- α 6 (KPNA5) |
| - CUL4A | - FBXO28 | - Ino80 |
| - CUL4B | - FBXO3 | - Inositol 1,4,5-trisphosphate receptor 3 |
| - Cup | - FBXO34 | - Integrin α 2 |
| - Cx43 | - FBXO38 | - Integrin α 4 |
| - cyclin B | - FBXO4 | - Integrin α L |
| - cyclophilin D (mitochondrial) | - FBXO40 | - Integrin α M β 2 |
| - CYP2D6 | - FBXO6 | |

- Integrin $\alpha V\beta 5$
- IL-1 β
- IRS-2
- Japanese encephalitis virus E protein
- JlpA
- KAP1
- KAT5
- KAT8
- KBTBD4
- KBTBD7
- KCBP
- KCNA5
- KCNA6
- KCNG1
- KCNS3
- KCNQ4
- KCTD8
- KDM3A/JMJD1A
- KDM4B/JMJD2B
- KEAP1
- KIAA0317
- Kir6.2
- KLHL1
- KLHL10
- KLHL13
- KLHL14
- KLHL15
- KLHL22
- KLHL23
- KLHL25
- KLHL26
- KLHL29
- KLHL32
- KLHL34
- KLHL36
- KLHL38
- KLHL6
- knob complexes (in the membrane of Plasmodium-infected erythrocytes)
- KSHV K1
- KSR1
- KSR2
- L protein of HRSV
- L protein of SHVV
- Lamin A/C
- LAMP-2A
- LANA of KS-HSV
- LAP
- LARP4B
- Legumain
- LGALS3BP
- LIS1
- LNX1
- LOC440248
- LOX1 (OLR1)
- LOXL2
- Lpl1 (*S. aureus*)
- LRP1 (=CD91)
- LRP5
- LRSAM1
- LSD1
- LSM8
- macromolecular aminoacyl-tRNA synthetase complex
- Macrophage scavenger receptor
- MAP1B
- MAP4
- MARCH9
- Mdm2
- MDM4
- MeCatalase1
- Mg²⁺-dependent phosphatidate phosphohydrolase
- MIF
- misfolded VHL
- MMP2, MMP3, MMP9
- μ -opioid receptor
- MRE11/Rad50/NBS1 (MRN) complex
- MRP1
- Msps/XMAP215/ch-TOG
- MTA1
- MTG8
- MUC1
- Myoglobin
- N-myc downstream-regulated gene 1 (NRDG1)
- N-WASP
- Na⁺-K⁺-Cl⁻ cotransporter 1
- NadA
- NAP1
- NB-LRR proteins: RPM1 and RPS2, Nod1, Nod2, NALP2, NALP3, NALP4, NALP12, IPAF, RPP4
- NBR1
- NCC
- NCT
- NDRG1
- NDRG2
- NELF-E
- Nervous necrosis virus capsid protein
- Neuraminidase
- Neuropeptide Y
- NHE1
- NHLRC1
- Nibrin
- NleH1 and NleH2
- NMNAT2
- Norovirus capsid protein VP1
- Nox1, Nox2, Nox3, Nox5
- NS1
- Nsl1
- nsP3 and nsP4 of Chikungunya virus
- Nucleoprotein (NP) of MERS-CoV
- Nup62
- Nwd1
- OGT
- OsCERK1
- P protein (rabies virus)
- P1 (picornaviral capsid precursor protein P1)
- P2X₇ purinergic receptor
- p14ARF
- p17 (Avian reovirus)
- p300
- PARK2
- PARK7 (DJ-1)
- PARP1
- PB1 and PB2 subunits of influenza RNA pol.
- PBD2
- PCGF1
- PCGF3
- PCNA
- PDE6 α (PDE6C), β , γ
- Peli1
- perilipin
- PfCRT
- PIDD
- Piwi
- PIWIL2
- PLC γ
- PLN
- polysomal ribonuclease 1 (PMR1)
- Porcine deltacoronavirus proteins N, NS7, NSP10
- PPAT
- PRDM14
- PRMT5
- pro-Dcp1
- prolactin receptor
- prostacyclin synthase
- proteasome
- PRPF8
- PRPF19
- PsbO
- PTPN22
- Ptx
- PUS7
- R-protein I-2
- R2TP complex through Pih1
- Rab- α GDI
- Rab3a
- Rab11a
- RAB40A
- Rac/Rop GTPase Rac1 (rice)
- Rac1
- Rad51
- Rad52
- RAG1
- Ral-binding protein 1 (RalBP1)
- RanBP9
- Rapsyn

- | | | |
|--|---|---|
| - Raptor | - SUR1 (subunit of β -cell ATP-sensitive potassium channel) | - USP14 |
| - RCBTB1 | - survivin | - Utp21 |
| - RCBTB2 | - SUV39H1 | - Vaccinia core protein 4a |
| - RCP | - SV40 large T-antigen | - vFLIP (of KSHV) |
| - Reovirus protein σ 1 | - Swr1 | - Vimentin |
| - REV1 | - α -synuclein | - VIP1 |
| - reverse transcriptase of hepatitis B virus | - Tab2/3 | - VP5 of pseudorabies virus |
| - RFWD3 | - Tas3 | - VP5, VP7, VP35 of grass carp reovirus |
| - RGS11 | - Tau protein | - VPS18 |
| - RGS6 | - Tax | - VPS41 |
| - RGS7 | - TCL1A | - WASF3 |
| - RGS9 | - telomerase | - WSB2 |
| - RhoB | - TFR1 | - WTAP |
| - RHOBTB1 | - thiopurine S-methyltransferase | - WWP1 |
| - ribosomal protein L2 (E. coli) | - thrombin receptor (PAR-1) | - XPO1 |
| - ribosomal proteins S3 and S6 | - thromboxane synthase | - XPORT |
| - ricin catalytic A chain | - TiIs | - XRCC1 |
| - RIG-I | - TIR1 | - ZEITLUPE |
| - RNA-dep. RNA polymerase (of Bamboo mosaic virus) | - Tissue plasminogen activator (tPA) | |
| - RNF10 | - Titin | |
| - RNF111 | - TLR4/MD-2 complex | |
| - RNF19B | - TLR7 | |
| - RNF40 | - TLR9 | |
| - RNGTT | - Tm-2 ² | |
| - Rnr4 | - TNFAIP3 (=A20) | |
| - Rpb1 | - TOM40 | |
| - RPN6 | - TRIM2 | |
| - Rsc3 | - TRIM7 | |
| - SA- β -galactosidase (GLB1) | - TRIM8 | |
| - SARS-CoV-2 proteins N, M, Orf3, Orf7a, and Orf7b | - TRIM10 | |
| - SCAP | - TRIM17 | |
| - SDCBP | - TRIM21 | |
| - SDF2 | - TRIM36 | |
| - SENP3 | - TRIM37 | |
| - SERCA2a | - TRIM41 | |
| - SERT (SLC6A4) | - TRIM49 | |
| - SF3B3 | - TRIM56 | |
| - SH3RF2 | - TRIM73 | |
| - Sicily | - TRIM74 | |
| - SIR2 (SIR2RP1 in Leishmania) | - Triosephosphate isomerase | |
| - SIRT1 | - Trithorax (and ortholog MLL) | |
| - SIRT2 | - Trx1 | |
| - SKP2 | - TrxR | |
| - SKP2 complexes | - TSG101 | |
| - SLC6A14 | - Tyrosine hydroxylase | |
| - SMURF1 | - UBA52 | |
| - SMYD1, SMYD2, SMYD3 | - UCH-L1 | |
| - snoRNP complexes | - UHRF1 | |
| - SNRNP200 | - UL42 (of HSV-1) | |
| - SOCS6 | - Ulp1 | |
| - SPSB1 | - uPA | |
| - SPSB3 | - Ura2 | |
| - SREC-I | - URI complex | |
| - STING | - Uroporphyrinogen decarboxylase (Heme) [in cyanobacteria] | |
| | - Us11 (of HSV-1) | |

Notes:

- Only the cytosolic form(s) of Hsp90 is considered.
- Clients from different species are all mixed together and the protein names are typically those of the original publication (i.e. not necessarily the official protein name).
- Only proteins are listed for which biochemical evidence for an interaction is available (i.e. geldanamycin effects alone are not considered as sufficient).
- more candidate interactors can be found in reports about proteomic approaches (Falsone et al. [2005] FEBS Lett. 579, 6350; Te et al. [2007] J. Proteome Res. 6, 1963; Caldas-Lopes et al. [2009] PNAS 106, 8368; Tsaytler et al. [2009] Cell Stress Chaperones 14, 629; Gong et al. [2009] Mol. Syst. Biol. 5, 275; Gano and Simon [2010] Mol. Cell. Proteomics 9, 255; Behrends et al. [2010] Nature 466, 68; Wang et al. [2010] Cancer Invest. 28, 635; Garcia-Descalzo et al. [2011] Cell Stress Chaperones 16, 203; Skarra et al. [2011] Proteomics 11, 1508, Moulik et al. [2011] Nat. Chem. Biol. 7, 818; Wu et al. [2012] Mol. Cell. Proteomics 11, M111 016675; Taipale et al. [2012] Cell 150, 987; Buljan et al. [2020] Mol. Cell 79, 504; Taipale et al. [2014] Cell 158, 434; Truman et al. [2015] J. Proteomics 112, 285; Savitsky et al. [2018] Cell 173, 260; Zhao et al. [2021] Mol. Cell 81, 2914; Liu et al. [2022] Nucleic Acids Res. 50, 6990), global analyses (e.g. Zhao et al. [2005] Cell 120, 715; Millson et al. [2005] Euk. Cell 4, 849; McClellan et al. [2007] Cell 131, 121; Franzosa et al. [2011] PLoS One 6, e28211; Sharma et al. [2012] Mol. Cell. Proteomics 11, M111 014654; Rizzolo et al. [2017] Cell Rep. 20, 2735; Miao et al. [2018] Anal. Chem. 90, 11751; O'Meara et al. [2019] PLoS Biol. 17, e3000358; Tsvetkov et al. [2020] Cell Rep. 32, 108001; Yan et al. [2023] Plant Cell Environ. 46, 1935; Kolhe et al. [2023] Mol. Cell 83, 2035); and in pharmacological surveys of kinases (Citri et al. [2006] J. Biol. Chem. 281, 14361; Haupt et al. [2012] BMC Cancer 12, 38).
- See **Hsp90Int.db** for the comprehensive (notably human) interactome built with data from public protein-protein interaction databases and the literature (Echeverría et al. [2011] PLoS One 6, e26044; and its associated database at <https://www.picard.ch/Hsp90Int>). Hsp90Int.db also uses exclusively the official NCBI names.
- Looking for references? See <https://www.picard.ch/downloads/Hsp90facts.pdf>.